

K_{e4} decays and Wigner cusp

Lucia Masetti

Johannes Gutenberg-Universität Mainz, Germany



**International Conference
“Heavy Quarks and Leptons”
Munich, 17th October 2006**



On behalf of the NA48/2 collaboration

Cambridge, CERN, Chicago, Dubna, Edinburgh, Ferrara, Firenze, Mainz,
Northwestern, Perugia, Pisa, Saclay, Siegen, Torino, Wien

1 Introduction

- Physics motivation
- The NA48/2 experiment: beamline and detector

2 $K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu_e$

- Form factors and $\pi\pi$ scattering lengths

3 $K^\pm \rightarrow \pi^0 \pi^0 e^\pm \nu_e$

- Branching fraction and form factors

4 $K^\pm \rightarrow \pi^0 \pi^0 \pi^\pm$

- The “cusp” effect
- Slopes and $\pi\pi$ scattering lengths

5 Conclusions

WHY?


- Very precise theoretical predictions from χPT (2%) and generalised χPT , depending only on **one free parameter**: **the quark condensate** $\langle \bar{q}q \rangle_0$
- $\langle \bar{q}q \rangle_0$ must be determined **experimentally**
- The size of $\langle \bar{q}q \rangle_0$ determines the **order** at which mass terms appear in the perturbative expansion

HOW?

- K_{e4} ($K \rightarrow \pi\pi e\nu$): no other hadrons, pions **close to threshold**
No theoretical uncertainty on the form factors, only on $a_0^2 = f(a_0^0)$
- $K^\pm \rightarrow \pi^0\pi^0\pi^\pm$: “cusp” at $M_{00}^2 = 4m_{\pi^+}^2$ due to **rescattering**
Theoretical uncertainty of the Cabibbo-Isidori model: **5%**

The NA48/2 beamline

Simultaneous K^+ and K^- beams with $p_K = (60 \pm 3) \text{ GeV}/c$
to measure charge asymmetry in $K \rightarrow 3\pi$ decays



The NA48/2 detector

- **Magnetic spectrometer:**

4 drift chambers

+ 1 dipole magnet

$\sigma(p)/p \simeq 0.9\%$ @ 20 GeV/


$(p_{\perp}^{kick} \simeq 120 \text{ MeV}/c)$

- **Electromagnetic calorimeter:**

Liquid krypton calorimeter

$\sigma(E)/E \simeq 1\%$ @ 20 GeV

$\sigma(t) \simeq 265 \text{ ps}$ for 50 GeV e^{-}





$K^\pm \rightarrow \pi^+\pi^-e^\pm\nu_e$: Selection criteria and reconstruction

- 3 tracks building a good vertex
- 1 electron with the same charge as the vertex (E/p and Linear Discriminant Analysis)
- Missing p_T : elliptical cut in the $(p_T, M_{3\pi})$ plane

Two reconstruction strategies:

- 1 Assume kaon mass, extract kaon momentum (quadratic equation)
- 2 Assume kaon momentum, extract kaon mass (linear equation)

Final sample: $\sim 370,000$ selected events



$K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu_e$: Background estimate

Expectation from MC simulation:

Channel	Rejection	Level
$K^\pm \rightarrow \pi^+ \pi^- \pi^\pm$ with $\pi \rightarrow e\nu$	Vertex selection	$\sim 0.2\%$
$K^\pm \rightarrow \pi^+ \pi^- \pi^\pm$ with π mis-ID	LDA, ellipse (p_T)	$\sim 0.2\%$
$K^\pm \rightarrow \pi^\pm \pi_D^0$ and $K^\pm \rightarrow \pi_D^0 \pi^0 \pi^\pm$	E/p for pion-ID	$< 0.01\%$

Estimate from DATA:


“Wrong sign” events ($\pi^\pm \pi^\pm e^\mp \nu_e$) can only be background ($\Delta S = \Delta Q$ rule)

Factor 2 scaling applied to background from $K^\pm \rightarrow \pi^+ \pi^- \pi^\pm$

Estimated background: $\sim 0.5\%$ of the signal, in agreement with the expectation


$K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu_e$: Form factor parametrisation

Kinematic variables



- $M_{\pi\pi}$
- $M_{e\nu}$
- $\cos \theta_\pi$ (in the $\pi\pi$ cm system)
- $\cos \theta_e$ (in the $e\nu$ cm system)
- ϕ (angle between the $\pi\pi$ and $e\nu$ planes in the K cm system)

Matrix element



$$T = \frac{G_F}{\sqrt{2}} V_{us}^* \bar{u}(p_\nu) \gamma_\mu (1 - \gamma_5) v(p_e) (V^\mu - A^\mu)$$





Form factors

$$\begin{aligned} F &= F_S e^{i\delta_0^0} + F_P e^{i\delta_1^1} \cos \theta_\pi \\ G &= G_P e^{i\delta_1^1} \\ H &= H_P e^{i\delta_1^1} \end{aligned}$$

$K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu_e$: Fit strategy

- A grid of **$10 \times 5 \times 5 \times 5 \times 12 = 15000$ equal population bins** was defined in the 5-dimensional space of the kinematic variables, separately for K^+ and K^-
- In the first step **10 independent 5 parameter** ($F_s, F_\rho, G_\rho, H_\rho, \delta$) **fits** were performed for each bin in $M_{\pi\pi}$, minimising a log-likelihood estimator that takes into account also the limited MC statistics
- In the second step the 5 sets of points were fitted with a **polynomial in powers of $q^2 = \frac{s}{4m_\pi^2} - 1$** , truncating the expansion according to the sensitivity
- The **dependence on $M_{e\nu}$** was found to be **negligible** within the total uncertainty and a possible f_e term was not included in the fit
- The **D-wave contribution** was also found to be **negligible** within the statistical uncertainty and was excluded from the fit


$K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu_e$: F, G and H









$$K^\pm \rightarrow \pi^+\pi^-\nu_e: a_0^0 \text{ and } a_0^2$$

The $\delta = \delta_0^0 - \delta_1^1$ distribution was fitted with a **1 parameter (a_0^0) function** given by the numerical solution of the Roy equations in Phys. Rept. 353, 207

a_0^0 and a_0^2 were constrained to lie on the **centre of the universal band**



$K^\pm \rightarrow \pi^+ \pi^- e^\pm \nu_e$: Data/MC comparison after the fit



$K^\pm \rightarrow \pi^+\pi^-\ e^\pm \nu_e$: Systematics and preliminary result

Systematic checks

- Two independent analyses with different **reconstruction methods**, **acceptance corrections**, **fit methods** and **MC parameters**
- Stability of **acceptance vs time** (variation of the simulated beam conditions)
- Uncertainty on **background estimate** (checked with data and MC)
- Uncertainty on **electron-ID efficiency** (variation of LDA cut)
- Uncertainty on **radiative corrections** (fraction of total effect with or w/o PHOTOS)
- Bias from **neglected s_e dependence** (MC tests)

NA48/2 preliminary result (2003 data)

$$f'_s/f_s = 0.169 \pm 0.009_{stat} \pm 0.034_{syst}$$

$$f''_s/f_s = -0.091 \pm 0.009_{stat} \pm 0.031_{syst}$$

$$f_p/f_s = -0.047 \pm 0.006_{stat} \pm 0.008_{syst}$$

$$g_p/f_s = 0.891 \pm 0.019_{stat} \pm 0.020_{syst}$$

$$g'_p/f_s = 0.111 \pm 0.031_{stat} \pm 0.032_{syst}$$

$$h_p/f_s = -0.411 \pm 0.027_{stat} \pm 0.038_{syst}$$

$$a_0^0 = 0.256 \pm 0.008_{stat} \pm 0.007_{syst}$$

$$\pm 0.018_{theor}$$

$K^\pm \rightarrow \pi^0 \pi^0 e^\pm \nu_e$: Selection and background estimate

- 4 photons compatible with two π^0 at the same vertex
- 1 electron track (E/p and shower width)
- Missing p_T : elliptical cut in the $(p_T, M_{3\pi})$ plane
- Using kaon momentum of 60 GeV/c along z, compute kaon mass (linear equation)

Background


Estimated from data
(reversing cuts):

$$K^\pm \rightarrow \pi^0 \pi^0 \pi^\pm$$

with pion mis-ID

$$K^\pm \rightarrow \pi^0 e^\pm \nu_e \gamma + \text{accidental photon}$$

Total contamination: $\sim 3\%$



$K^\pm \rightarrow \pi^0 \pi^0 e^\pm \nu_e$: Branching ratio

Selected events

K_{e4}^{00} candidates: 9642, with a **background** contamination of 276 ± 94

Normalisation ($K^\pm \rightarrow \pi^0 \pi^0 \pi^\pm$) $\sim 8 \times 10^6$, negligible background

Systematic uncertainties

Acceptance, trigger efficiency and energy measurement of the calorimeter

NA48/2 preliminary result (2003 data)

$$BR(K_{e4}^{00}) = (2.587 \pm 0.026_{stat} \pm 0.019_{syst} \pm 0.029_{ext}) \times 10^{-5}$$

$K^\pm \rightarrow \pi^0 \pi^0 e^\pm \nu_e$: Form factors



Same formalism as for K_{e4}^{+-} , but, for the symmetry of the $\pi^0 \pi^0$ system,
no P-wave!

Effect of cusp at $M_{00}^2 = 4m_{\pi^+}^2$ taken into account, 37,700 events used

NA48/2 preliminary result (2003+2004 data)

$$f'_s/f_s = 0.129 \pm 0.036_{stat} \pm 0.020_{syst}$$

$$f''_s/f_s = -0.040 \pm 0.034_{stat} \pm 0.020_{syst}$$





$K^\pm \rightarrow \pi^0 \pi^0 \pi^\pm$: Cusp in the M_{00}^2 distribution

Selected events (2003): 23×10^6

M_{00}^2 computed imposing the mean vertex of the π^0 's
(improved resolution close to threshold)

Evidence for a cusp at $M_{00}^2 = 4m_{\pi^+}^2$ due to $\pi\pi$ rescattering



Fit to the Cabibbo-Isidori model (JHEP03 021):

$$M_0 = A_0 \left(1 + \frac{1}{2} g_0 u + \frac{1}{2} h' u^2 \right)$$

Phys. Lett. B 633 (2006) 173

$$\begin{aligned} g_0 &= 0.645 \pm 0.004_{stat} \pm 0.009_{syst} \\ h' &= -0.047 \pm 0.012_{stat} \pm 0.011_{syst} \\ a_2 &= -0.041 \pm 0.022_{stat} \pm 0.014_{syst} \\ a_0 - a_2 &= 0.268 \pm 0.010_{stat} \pm 0.004_{syst} \\ &\quad \pm 0.013_{theor} \end{aligned}$$

$K^\pm \rightarrow \pi^0 \pi^0 \pi^\pm$: $k'v^2$ term


Fit in two steps

- 1 Fit the modified matrix element

$$M_0 = A_0 \left(1 + \frac{1}{2} g_0 u + \frac{1}{2} h' u^2 + \frac{1}{2} k' v^2 \right)$$

above the cusp in the plane $\cos \theta$ vs M_{00}^2 $\cos \theta =$ angle between π^+ and π^0 in $\pi^0 \pi^0$ cm

- 2 Reweight the MC with the obtained value of k' and fit the M_{00}^2 distribution with the **Cabibbo-Isidori model** to obtain the cusp parameters



Systematic checks:

Acceptance, trigger efficiency


NA48/2 preliminary result (2003 data)

$$k' = 0.0097 \pm 0.0003_{stat} \pm 0.0008_{syst}$$

Note: h' and k' are not the same as the PDG parameters



Comparison of the scattering length results



New results **compatible** with theory and previous measurements

K_{e4} : **higher sensitivity** than BNL E865 (higher statistics at high $M_{\pi\pi}$)

Conclusions

- Form factor and $\pi\pi$ scattering length measurement from K_{e4}^{+-}

$$a_0^0 = 0.256 \pm 0.008_{stat} \pm 0.007_{syst} \pm 0.018_{theor}$$

Compatible with χPT prediction and previous results, higher sensitivity

- Branching ratio and form factor measurements from K_{e4}^{00}

$$BR(K_{e4}^{00}) = (2.587 \pm 0.026_{stat} \pm 0.019_{syst} \pm 0.029_{ext}) \times 10^{-5}$$

BR: factor 8 better than latest measurement

Form factors: compatible with charged channel

- $\pi\pi$ scattering length and slope measurements from $K^\pm \rightarrow \pi^0\pi^0\pi^\pm$

$$a_0 - a_2 = 0.268 \pm 0.010_{stat} \pm 0.004_{syst} \pm 0.013_{theor}$$

- First evidence for a value of $k \neq 0$ in the $K^\pm \rightarrow \pi^0\pi^0\pi^\pm$ Dalitz plot

$$k' = 0.0097 \pm 0.0003_{stat} \pm 0.0008_{syst}$$