

Measurement of FCNC decays $K^\pm \rightarrow \pi^\pm l^+ l^-$ by NA48/2 at CERN

Evgueni Goudzovski
(University of Birmingham)

on behalf of the **NA48/2** collaboration

(Cambridge, CERN, Chicago, Dubna, Edinburgh, Ferrara, Florence, Mainz,
Northwestern, Perugia, Pisa, Saclay, Siegen, Turin, Vienna)

Outline:

- 1) Beams, detector and data taking in 2003/04;
- 2) $K^\pm \rightarrow \pi^\pm e^+ e^-$ analysis: NA48/2 final results;
- 3) $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ analysis: analysis status and prospects;
- 4) Summary.

NA48/NA62 experiments

1997: ε'/ε run K_L+K_S

1998: K_L+K_S

1999: K_L+K_S

K_S HI

2000: K_L only

K_S HI

2001: K_L+K_S

K_S HI

2002: K_S /hyperons HI

2003: K^+/K^-

2004: K^+/K^-

2007: $K_{e^2}^{\pm}/K_{\mu^2}^{\pm}$

tests

2008: $K_{e^2}^{\pm}/K_{\mu^2}^{\pm}$

tests

2006–2010:
design & construction

2011:
start of $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ run

NA48/NA62: a series of experiments,
present-day CERN kaon physics programme

The presented results are based on 2003/04 data

NA48

NA48/1


NA48/2

NA62

(phase I)

NA62

(phase II)



Jura mountains

France

NA48/NA62

SPS

LHC


Geneva airport

Switzerland

N


NA48/2: kaon beam line

Kaon momentum spectrum



2-3M K/spill ($\pi/K \sim 10$),
 π decay products stay in pipe.
 Flux ratio: $K^+/K^- \approx 1.8$


Simultaneous K^+ and K^- beams:
 large charge symmetrization of
 experimental conditions



Be target
 $\sim 7 \cdot 10^{11}$ ppp,
 400 GeV
 0.36 mrad

Beams coincide within ~ 1 mm
 all along 114m decay volume


- Front-end achromat**
 - Momentum selection
- Quadrupole quadruplet**
 - Focusing
 - μ sweeping
- Cleaning**
- Beam spectrometer (momentum resolution $\sim 0.7\%$)**



The NA48 detector

Main detector components:

- **Magnetic spectrometer (4 DCHs):**
4 views/DCH: redundancy \Rightarrow efficiency;
used in trigger logic;
 $\Delta p/p = 1.0\% + 0.044\% \cdot p$ [GeV/c].
- **Hodoscope**
fast trigger;
precise track time measurement (150ps).
- **Liquid Krypton EM calorimeter (LKr)**
High granularity, quasi-homogenous;
 $\sigma_E/E = 3.2\%/E^{1/2} + 9\%/E + 0.42\%$ [GeV];
 $\sigma_x = \sigma_y = 0.42/E^{1/2} + 0.6\text{mm}$ (1.5mm@10GeV).
Used for γ detection and particle ID.
- **Hadron calorimeter, muon veto counters,**
photon vetoes.



NA48/2 data taking: completed

A view of the NA48/2 beam line



2003 run: ~ 50 days

2004 run: ~ 60 days

$K_{3\pi}$ statistics in 2 years:

$K^{\pm} \rightarrow \pi^{-}\pi^{+}\pi^{\pm}$: $\sim 4 \cdot 10^9$

$K^{\pm} \rightarrow \pi^0\pi^0\pi^{\pm}$: $\sim 1 \cdot 10^8$

Rare K^{\pm} decays:
BRs down to 10^{-9}
can be measured

>200 TB of data recorded

$K_{\pi\parallel}$: motivation & theory

$K^\pm \rightarrow \pi^\pm \gamma^* \rightarrow \pi^\pm l^+ l^-$: suppressed FCNC process proceeding through single virtual photon exchange. Information on weak interactions at low energy.

$$d\Gamma_{\pi ee}/dz \sim P(z) \cdot |W(z)|^2 \quad z=(M_{ee}/M_K)^2, P(z) \text{ is a phase space factor}$$

Considered models for the form factor:

- (1) polynomial: $W(z) = G_F M_K^2 \cdot f_0 \cdot (1 + \delta z)$
- (2) ChPT $O(p^6)$: $W(z) = G_F M_K^2 \cdot (a_+ + b_+ z) + W^{\pi\pi}(z)$
G. D'Ambrosio et al., JHEP 9808 (1998) 4
- (3) ChPT, large- N_c QCD: $W(z) = W(w, \beta, z)$
S. Friot, D. Greynat, E. de Rafael, PLB 595 (2004) 301
- (4) "Mesonic" ChPT: $W(z) = W(M_a, M_\rho, z)$
A.Z. Dubnickova et al., Phys. Part. Nucl. Lett. 5 (2008) 76 [hep-ph/0611175]

- Goals:
- 1) $d\Gamma/dz$ and model-independent BR in kinematic range $z > 0.08$;
 - 2) parameters of the models + BRs in the full kinematic range;
 - 3) upper limit for CPV charge asymmetry of decay rates.

Principal selection criteria

The $K^{\pm} \rightarrow \pi^{\pm} e^{+} e^{-}$ is measured normalizing to $K^{\pm} \rightarrow \pi^{\pm} \pi^{0} D \rightarrow \pi^{\pm} e^{+} e^{-} \gamma$.
Thus particle ID efficiencies cancel in the first order.

Common selection criteria:

3-track vertex [consistent in space/time],

one π candidate, two opposite sign electron candidates.

Electron (pion) ID based on E deposition : $E/p > 0.95$ ($E/p < 0.85$).



Signal selection:

Kinematic suppression of $\pi^{\pm} \pi^{0} D$
background: $M_{ee} > 140 \text{ MeV}/c^2$.

Limitations on reconstructed
 $\pi^{\pm} e^{+} e^{-}$ invariant mass,
total & transverse momentum




Normalization selection:

Selection of good γ candidate.
Limitations on reconstructed
 $e^{+} e^{-} \gamma$ and $\pi^{\pm} e^{+} e^{-} \gamma$ masses,
total & transverse momentum.

Signal & normalisation samples


Total kaon decays in fiducial volume (2003+2004): $N_K = 1.7 \times 10^{11}$



7253 candidates

Background/Signal = 1.0%

($K^\pm \rightarrow \pi^\pm \pi^0_D$, $K^\pm \rightarrow \pi^0_D e^\pm \nu$ + particle misID, two $e^+ e^-$ pairs: $\pi^0_{D(D)}$, γ conversions).
Subtracted with same-sign lepton candidates.




12.12 mln candidates

Background/Signal = 0.15%

($K^\pm \rightarrow \pi^0_D \mu^\pm \nu$, subtracted with simulation)


The accessible M_{ee} region

All analysis cuts except the M_{ee} cut are applied




- The region $M_{ee} < 140 \text{ MeV}/c^2$ is dominated by background and not accessible;
- Subtraction of the $K^\pm \rightarrow \pi^\pm \pi^0_{DD}$ reveals the $\pi^0 \rightarrow e^+ e^-$ signal (~ 500 events).

Kinematic variables: $(z, \cos\theta_{e\pi})$




z distribution is sensitive to the form-factor and contains the dynamical information:
 $d\Gamma/dz \sim P(z) \times |W(z)|^2$.



Decay via one photon exchange:
 $d\Gamma/d\theta \sim \sin^2\theta = (1 - \cos^2\theta)$,
 θ = angle between (e^+, π) in (e^+, e^-) frame.
 No dynamical information in this projection.

Fit results (1)




Model parameters with their statistical errors and correlation coefficients:

$$\begin{aligned}
 (1) \left\{ \begin{array}{l} f_0 = 0.531 \pm 0.012 \\ \delta = 2.32 \pm 0.15 \\ \rho(\delta, f_0) = -0.962 \end{array} \right. \\
 (2) \left\{ \begin{array}{l} a_+ = -0.578 \pm 0.012 \\ b_+ = -0.779 \pm 0.053 \\ \rho(a_+, b_+) = -0.913 \end{array} \right. \\
 (3) \left\{ \begin{array}{l} w = 0.057 \pm 0.005 \\ \beta = 3.45 \pm 0.24 \\ \rho(w, \beta) = 0.999 \end{array} \right. \\
 (4) \left\{ \begin{array}{l} M_a = (0.951 \pm 0.028) \text{ GeV} \\ M_\rho = (0.705 \pm 0.010) \text{ GeV} \\ \rho(M_a, M_\rho) = 0.998 \end{array} \right.
 \end{aligned}$$





Fit results (2)

Squared form factors
normalised by $|W(0)|^2=1$



Fit results (3)

68% CL contours



Fit results (4)

$BR_{mi} \times 10^7 =$	$2.28 \pm 0.03_{stat} \pm 0.04_{syst} \pm 0.06_{ext}$	$=$	2.28 ± 0.08
$f_0 =$	$0.531 \pm 0.012_{stat} \pm 0.008_{syst} \pm 0.007_{ext}$	$=$	0.531 ± 0.016
$\delta =$	$2.32 \pm 0.15_{stat} \pm 0.09_{syst}$	$=$	2.32 ± 0.18
$BR_1 \times 10^7 =$	$3.05 \pm 0.04_{stat} \pm 0.05_{syst} \pm 0.08_{ext}$	$=$	3.05 ± 0.10
$a_+ =$	$-0.578 \pm 0.012_{stat} \pm 0.008_{syst} \pm 0.007_{ext}$	$=$	-0.578 ± 0.016
$b_+ =$	$-0.779 \pm 0.053_{stat} \pm 0.036_{syst} \pm 0.017_{ext}$	$=$	-0.779 ± 0.066
$BR_2 \times 10^7 =$	$3.14 \pm 0.04_{stat} \pm 0.05_{syst} \pm 0.08_{ext}$	$=$	3.14 ± 0.10
$w =$	$0.057 \pm 0.005_{stat} \pm 0.004_{syst} \pm 0.001_{ext}$	$=$	0.057 ± 0.007
$\beta =$	$3.45 \pm 0.24_{stat} \pm 0.17_{syst} \pm 0.05_{ext}$	$=$	3.45 ± 0.30
$BR_3 \times 10^7 =$	$3.13 \pm 0.04_{stat} \pm 0.05_{syst} \pm 0.08_{ext}$	$=$	3.13 ± 0.10
$M_a =$	$0.974 \pm 0.030_{stat} \pm 0.019_{syst} \pm 0.002_{ext}$	$=$	0.974 ± 0.035 [GeV/c]
$M_\rho =$	$0.716 \pm 0.011_{stat} \pm 0.007_{syst} \pm 0.002_{ext}$	$=$	0.716 ± 0.014 [GeV/c]
$BR_4 \times 10^7 =$	$3.18 \pm 0.04_{stat} \pm 0.05_{syst} \pm 0.08_{ext}$	$=$	3.18 ± 0.10

Including uncertainty due to the model dependence,


$$BR = (3.11 \pm 0.04_{stat} \pm 0.05_{syst} \pm 0.08_{ext} \pm 0.07_{model}) \times 10^{-7} = (3.11 \pm 0.12) \times 10^{-7}$$

CPV parameter (first measurement; only uncorrelated K^+/K^- uncertainties):

$$\Delta(K_{\pi ee}^\pm) = (BR^+ - BR^-) / (BR^+ + BR^-) = (-2.1 \pm 1.5_{stat} \pm 0.6_{syst}) \times 10^{-2}$$

Comparison: FF slope δ


Measurement	Process	Result
Alliegro et al., PRL 68 (1992) 278	$K^+ \rightarrow \pi^+ e^+ e^-$	1.31 ± 0.48
Appel et al. [E865], PRL 83 (1999) 4482	$K^+ \rightarrow \pi^+ e^+ e^-$	2.14 ± 0.20
Ma et al. [E865], PRL 84 (2000) 2580	$K^+ \rightarrow \pi^+ \mu^+ \mu^-$	$2.45^{+1.30}_{-0.95}$
NA48/2, arXiv:0903:3130 (2009)	$K^\pm \rightarrow \pi^\pm e^+ e^-$	2.32 ± 0.18



- NA48/2 measurement of δ is compatible with the earlier results, has good precision;
- A contradiction of the data to the meson dominance models observed earlier is further confirmed;
- NA48/2 values of $(f_0, a_+, b_+, w, \beta)$ are in agreement with BNL E865 ones.

Comparison: BR in full z range


Measurement	Sample	BR×10 ⁷
Bloch et al., PL 56 (1975) B201	41 (K ⁺)	2.70±0.50
Alliegro et al., PRL 68 (1992) 278	500 (K ⁺)	2.75±0.26
Appel et al. [E865], PRL 83 (1999) 4482	10,300 (K ⁺)	2.94±0.15
NA48/2, arXiv:0903:3130 (2009)	7,300 (K [±])	3.11±0.12



Comparison of E865 vs NA48/2 results taking into account correlated uncertainties (normalization and model dependence):
 1.6 σ difference.


The NA48/2 results are final and published:
 CERN-PH-EP-2009-005, arXiv:0903:3130,
 accepted by Physics Letters B.

$K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ analysis (1)



NA48/2: $N=3,100$ $K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ candidates with 3% background.
Background well described by both $K_{3\pi}$ MC and same-sign muon samples.
Cf. total world sample is ~ 700 events.

$K^\pm \rightarrow \pi^\pm \mu^+ \mu^-$ analysis (2)



Analysis is well advanced,
we aim to present the preliminary results in 2009.

- Precise study of the $K^{\pm} \rightarrow \pi^{\pm} e^{+} e^{-}$ decay ($BR \sim 3 \times 10^{-7}$):
 - sample & precision comparable to world's best ones;
 - (BR, FF) agree to theory and earlier measurements;
 - first limit on the CPV charge asymmetry obtained;
 - final results published recently.
- Precise study of the $K^{\pm} \rightarrow \pi^{\pm} \mu^{+} \mu^{-}$ decay ($BR \sim 0.8 \times 10^{-7}$):
 - NA48/2 sample ~ 4 times larger than total world sample;
 - background is low and under control;
 - aim at preliminary results in 2009 – stay tuned!